

## ***Gingeras Lab RNA-Seq Library Production Document***

### **ENCODE Transcriptome**

Sample Description: HsAVEC 093WC Small TAP

RNA ID: 093WC

Library ID: LID47615

Protocol ID:

---

### **Cold Spring Harbor Laboratory**

Genome Center

500 Sunnyside Blvd

Woodbury, NY 11797

### **LAB MEMBERS**

Wet lab: Meagan Fastuca, Carrie A. Davis, Jorg Drenkow, Lei Hoon See, Huaiyen Wang.

Computational Lab: Alex Dobin, Sonali Jha, Wei Lin, Felix Schlesinger, Chris Zaleski, Chenghai Xue.

PI: Tom Gingeras

**STRATEGY:** This document contains information about small RNA libraries generated as part of the ENCODE Consortia. It describes the Small RNA Isolation and cloning methods used to generate stranded libraries that capture the 5' ends of RNAs <200 nucleotides in length. The libraries can then be sequenced on the Illumina platform. The 3' ends are A-tailed followed by ligating on a RNA linker to the 5' ends and RT-PCR.

### PromoCell RNA ISOLATION:

#### Kits: mirVana miRNA Isolation Kit (Cat #: AM1560)

1. Thaw the cells which are in the RNAlater Buffer. Transfer the cells and RNAlater buffer to a new 15ml Falcon RNase free tube. Add 6 times volume Lysis/Binding Solution to the tube, mix them well by vortex.
2. Homogenize the sample using a syringe and 18 gauge needle. Pass sample through the needle about twenty times.
3. Add 1/10 volume of miRNA Homogenate Additive to the homogenate, and mix well by vortexing or inverting the tube several times.
4. Leave the mixture on ice for 10 min.
5. Add a volume of Acid-Phenol:Chloroform that is equal to the lysate volume before addition of the miRNA Homogenate Additive. For example, if the original lysate volume was 300  $\mu$ L, add 300  $\mu$ L Acid-Phenol:Chloroform. (Be sure to withdraw from the bottom phase in the bottle of Acid-Phenol:Chloroform, because the upper phase consists of an aqueous buffer.)
6. Vortex for 30–60 sec to mix. Leave at room temperature for 2 minutes.
7. Centrifuge for 5 min at maximum speed (10,000 x g) at room temperature to separate the aqueous and organic phases. After centrifugation, the interphase should be compact; if it is not, repeat the centrifugation.
8. Carefully remove the aqueous (upper) phase without disturbing the lower phase, and transfer it to a fresh tube (DO NOT DISCARD). Note the volume removed.
- 9.

#### Separating total RNA Procedure

1. Preheat Elution Solution to 95°C for use in eluting the RNA from the filter at the end of the procedure. If the 100% ethanol you plan to use for this procedure is stored cold, warm it to room temperature before starting the Final RNA Isolation.
2. Add 1.25 volume of 100% ethanol to the aqueous phase recovered from the organic extraction. Mix thoroughly by inverting the tube several times.
3. For each sample, place a Filter Cartridge into one of the Collection Tubes supplied. Pass the sample through a Filter Cartridge, and collect the filtrate. Up to 700  $\mu$ L can be applied to a Filter Cartridge at a time. For sample volumes greater than 700  $\mu$ L, apply the mixture in successive applications to the same filter.
4. Centrifuge for ~15 sec to pass the mixture through the filter. Centrifuge at RCF 10,000 x g (typically 10,000 rpm). Spinning harder than this may damage the filters.
5. Apply 700  $\mu$ L miRNA Wash Solution 1 (working solution mixed with ethanol) to the Filter Cartridge from above and centrifuge for ~5–10 sec or use a vacuum to pull the solution through the filter. Discard the flow-through from the Collection Tube, and replace the Filter Cartridge into the same Collection Tube.
6. Apply 500  $\mu$ L Wash Solution 2/3 (working solution mixed with ethanol) and draw it through the Filter Cartridge as in the previous step.
7. Repeat with a second 500  $\mu$ L aliquot of Wash Solution 2/3.
8. After discarding the flow-through from the last wash, replace the Filter Cartridge in the same Collection Tube and spin the assembly for 1 min to remove residual fluid from the filter.
9. Transfer the Filter Cartridge into a fresh Collection Tube (provided with the kit). Apply 100  $\mu$ L of pre-heated (95°C) nuclease-free water to the center of the filter, and close the cap. Spin for ~20–30 sec at maximum speed to recover the RNA.
10. Transfer the RNA solution to a new RNase free eppendorf tube. Follow by the Separating large RNA procedure.
- 11.

#### Separating the small RNA Procedure

1. Mix total RNA with 5 volumes Lysis/Binding Buffer.
2. Add 1/10 volume of miRNA Homogenate Additive to the RNA mixture from the previous step, and mix well by vortexing or inverting the tube several times. Leave the mixture on ice for 10 min.
3. Add 1/3 volume of 100% ethanol to the RNA mixture from the previous step. Mix thoroughly by inverting the tube several times.
4. For each sample, place a Filter Cartridge into one of the Collection Tubes supplied. Pipet the lysate/ethanol mixture (from the previous step) onto the Filter Cartridge. Up to 700  $\mu$ L can be applied to a Filter Cartridge at a time. For sample volumes greater than 700  $\mu$ L, apply the mixture in successive applications to the same filter. Centrifuge for 30 seconds to pass the mixture through the filter. Centrifuge at RCF 5,000  $\times$  g (typically 5,000 rpm). Collect the filtrate. If the RNA mixture is >700  $\mu$ L, transfer the *flow-through* to a fresh tube, and repeat until all of the RNA mixture is through the filter. Pool the collected filtrates if multiple passes were done, and measure the total volume of the filtrate. After discarding the flow-through from the last wash, replace the Filter Cartridge in the same Collection Tube and spin the assembly for 1 min to remove residual fluid from the filter.
5. Add 2/3 volume room temperature 100% ethanol to filtrate (i.e. flow-through).
6. For each sample, place a Filter Cartridge into one of the Collection Tubes supplied. Pipet the filtrate/ethanol mixture (from the previous step) onto a second Filter Cartridge. Up to 700  $\mu$ L can be applied to a Filter Cartridge at a time. For sample volumes greater than 700  $\mu$ L, apply the mixture in successive applications to the same filter. Centrifuge for ~1 min to pass the mixture through the filter. Centrifuge at RCF 5,000  $\times$  g (typically 5,000 rpm). Discard the flow-through, and repeat until all of the filtrate/ethanol mixture is through the filter. Reuse the Collection Tube for the washing steps.
7. Apply 700  $\mu$ L miRNA Wash Solution 1 (working solution mixed with ethanol) to the Filter Cartridge and centrifuge for ~1 min at RCF 5,000  $\times$  g, or use vacuum to pass the solution through the filter. Discard the flow-through from the Collection Tube, and replace the Filter Cartridge into the same Collection Tube.
8. Apply 500  $\mu$ L Wash Solution 2/3 (working solution mixed with ethanol) and draw it through the Filter Cartridge as in the previous step.
9. Repeat with a second 500  $\mu$ L aliquot of Wash Solution 2/3. After discarding the flow-through from the last wash, replace the Filter Cartridge in the same Collection Tube and spin the assembly for 1 min at RCF 10,000  $\times$  g to remove residual fluid from the filter.
10. Transfer the Filter Cartridge into a fresh Collection Tube (provided with the kit). Apply 50  $\mu$ L of 95°C Elution Solution, and close the cap. Incubate at room temperature for ~2 min. Spin for 1 min at RCF 10,000  $\times$  g to recover the RNA. Repeat steps 9 with a second aliquot of preheated Elution Solution.
11. Transfer the RNA solution to a new RNase free 1.5ml tube. Follow by Ethanol Precipitation.

### Ethanol Precipitation

1. Add 2.5 volumes of 100% ethanol and 1/10 volumes of NaOAc PH 5.5 (Ambion Cat. # AM9740) to the eluted RNA.
2. Freeze in -80°C for at least 30 min.
3. Centrifuge for 35 min at max speed at 4°C.
4. Pipette and discard the supernatant making sure not to touch the pellet of RNA.
5. Wash with 1 mL of 70% ethanol and centrifuge at max speed for 5 min.
6. Pipette and discard the supernatant.
7. Open the cap and speed vacuum at low heat for 3-5 min making sure that the pellet is dry.
8. Resuspend the pellet with RNase-free water.

### DNase Digest (same for Small and Large RNA)

## Clean up

**RIBOMINUS TREATMENT:** In addition to the probes supplied with the Ribominus kit we also spike-in our own LNA probes against the 5S and 5.8S rRNA.

5S-LNAprobe-1 tt+Ccc+Agg+Cgg+Tct+Ccc+At

5S-LNAprobe-2 tc+Agg+Gtg+Gta+Tgg+Ccg+Tag

5.8S-LNAprobe-1 ct+Tca+Tcg+Acg+Cac+Gag+Cc

5.8S-LNAprobe-2 cg+Ctc+Aga+Cag+Gcg+Tagc

## Hybridization Step

Instructions are provided below to perform hybridization for 1–10 µg of your total RNA sample with the RiboMinus™ Eukaryote Probe. To process >10 µg total RNA sample, divide your sample into two samples, each containing <10 µg total RNA.

1. Set a water bath or heat block to 70–75°C.

2. To a sterile, RNase-free 1.5 mL microcentrifuge tube, add the following:

Total RNA (1–10 µg): <10 µL

RiboMinus™ Probe (15 pmol/L): 8 µL

Custom probes (5S, 5.8S 100uM each) : 1.5 µL

Hybridization Buffer: 100 µL

3. Incubate the tube at 70–75°C for 5 minutes to denature RNA.

4. Allow the sample to cool to 37°C slowly over a period of 30 min by placing the tube in a 37°C water bath (a heat block works as well). To promote sequence-specific hybridization, it is important to allow slow cooling. **Do not** cool samples quickly by placing tubes in cold water.

5. While the sample is cooling down, proceed to **Preparing Beads**.

\*An earlier version of this protocol says to use RNA in less than 20 µL, add 10 µL of probe and 300 µL hybridization buffer, this larger volume means you need to precipitate the ribominused RNA in a 2 mL tube at the end. Either way works. It doesn't change anything else expect the supernatant volumes and the precipitation tube size.

## Preparing Beads

1. Resuspend RiboMinus™ Magnetic Beads in its bottle by thorough vortexing.

2. Pipet 750 µL of the bead suspension into a sterile, RNase-free, 1.5 mL microcentrifuge tube.

3. Place the tube with the bead suspension on a magnetic separator for 1 min. The beads settle to the tube side that faces the magnet. Gently aspirate and discard the supernatant.

4. Add 750 µL sterile, DEPC Water to the beads and resuspend beads by pipetting

5. Place tube on a magnetic separator for 1 min. Aspirate and discard the supernatant.

6. Repeat Steps 4–5 once.

7. Resuspend beads in 750 µL Hybridization Buffer and transfer 250 µL beads to a new tube and maintain the tube at 37°C for use at a later step.

8. Place the tube with 500 µL beads on a magnetic separator for 1 min. Aspirate and discard the supernatant.

9. Resuspend beads in 200 µL Hybridization Buffer and keep the beads at 37°C until use.

## Removing rRNA

1. After the incubation at 37°C for 30 min of the hybridized sample (above), briefly centrifuge the tube to collect the sample to the bottom of the tube.

2. Transfer the sample (~120 µL - this will be ~330 µL with the older protocol) to the prepared RiboMinus™ Magnetic beads from Step 9 (**Preparing Beads**, above). Mix well by pipetting up and down

3. Incubate the tube at 37°C for 15 min. During incubation, gently mix the contents occasionally. Briefly centrifuge the tube to collect the sample to the bottom of the tube.

4. Place the tube on a magnetic separator for 1 min to pellet the rRNA-probe complex. **Do not discard the supernatant. The supernatant contains RiboMinus™ RNA.**

5. Place the tube with 250  $\mu$ L beads from Step 7 (**Preparing Beads**, above) on a magnetic separator for 1 min. Aspirate and discard the supernatant.
6. To this tube of beads, add ~320  $\mu$ L (~500  $\mu$ L with older protocol) supernatant containing RiboMinus™ RNA from Step 4, above. Mix well by pipetting up and down or low speed vortexing.
7. Incubate the tube at 37°C for 15 min. During incubation, gently mix the contents occasionally. Briefly centrifuge the tube to collect the sample to the bottom of the tube.
8. Place the tube on a magnetic separator for 1 min to pellet the rRNA-probe complex. **Do not discard the supernatant as the supernatant contains RiboMinus™ RNA.**
9. Transfer the supernatant (~ 320  $\mu$ L - ~500  $\mu$ L with older protocol) containing **RiboMinus™ RNA** to a small filter column and spin at max speed for 2 min to remove any remaining magnetic particles.
10. Transfer flow through (ribominus RNA) to a new tube (1.5 mL for small volume, 2 mL for large volume)
11. Ethanol precipitate as before but add 1  $\mu$ L glycoblue to facilitate the precipitation.
12. After drying the pellet, resuspend in 21  $\mu$ L H<sub>2</sub>O. Keep 1  $\mu$ L for running a small RNA Bioanalyzer chip.

## LIBRARY PROTOCOL:

The following primers and RNA Linker are needed to perform this procedure:

5'SBS3\_Adapter (This is the RNA ligated onto the 5' end): "r" = ribose, RNA base  
**5'-rArCrArCrUrCrUrUrCrCrUrArCrArCrGrArCrGrCrUrCrUrUrCrCrGrArUrCrUrNrNrNrCrG**  
 A-Tail RT Primer (This is the primer used in the RT reaction):  
**5'-TCTCGGCATTCTCTGCTGAACCGCTCTTCGGATCTTTTTTTTTTTVN**  
 PE 5' PCR (PCR Primer):  
**5'-AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATC**  
 PE 3' PCR (PCR Primer):  
**5'-CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCTGCTGAACCGCTCTTC**

## Tobacco Acid Pyrophosphatase Reaction (TAP, Epicentre T19250)

1. Denature RNA at 85°C for 2 min or 65°C for 10 min. Cool on ice for 1 min.
2. Set up the reaction by adding :
 

a. RNA	21.25 $\mu$ L
b. 10X TAP reaction buffer	2.5 $\mu$ L
c. Anti-RNase (Ambion 20U/ $\mu$ L)	1 $\mu$ L
d. TAP (10 U/ $\mu$ L)	0.25 $\mu$ L
3. Incubate at 37°C for 1 hr.
4. Proceed to small RNA column cleanup, then ethanol precipitation.
5. After drying the pellet, resuspend in 28  $\mu$ L H<sub>2</sub>O.

## Small RNA Column Cleanup

1. Bring sample up to 100  $\mu$ L. (Make sure there is a max of 45  $\mu$ g RNA in the 100  $\mu$ L RNA sample.)
2. Add 350  $\mu$ L Buffer RLT to the 100  $\mu$ L sample of RNA. Vortex to mix well.
3. Add 675  $\mu$ L of 100% ethanol to the reaction and mix by inverting.
4. Transfer 700  $\mu$ L of sample into an RNeasy MinElute column in a 2 mL collection tube. Close the lid gently and centrifuge for 30 s at 8000 x g ( $\geq 10,000$  rpm). Discard the flow through. Repeat this step with the remaining sample.
5. Add 700  $\mu$ L Buffer RWT to the RNeasy MinElute spin column. Close the lid gently and centrifuge for 30 s at  $\geq 8000$  x g ( $\geq 10,000$  rpm) to wash the column. Discard the flow through.
6. Pipet 500  $\mu$ L Buffer RPE into the RNeasy MinElute spin column. Close the lid gently and centrifuge for 30 s at  $\geq 8000$  x g ( $\geq 10,000$  rpm). Discard the flow-through.
7. Add 500  $\mu$ L of 80% ethanol to the RNeasy MinElute spin column. Close the lid gently and centrifuge for 2 min at  $\geq 8000$  x g ( $\geq 10,000$  rpm) to dry the spin column membrane. Discard the flow-through and the collection tube.
8. Place the RNeasy MinElute spin column into a new 2 mL collection tube, making sure that the column does not come in contact with the flow through. Open the lid and centrifuge for 5 min at  $\geq 8000$  x g ( $\geq 10,000$  rpm).
9. Place the RNeasy MinElute spin column into a 1.5 mL collection tube and pipet 20  $\mu$ L RNase free water onto the spin column membrane. Close the lid gently and wait 1 min. Then centrifuge for 1 min at  $\geq 8000$  x g ( $\geq 10,000$  rpm).

10. Repeat step 9 with a second volume of 20  $\mu$ L RNase free water.
11. Proceed to ethanol precipitation.

#### A-tailing (PolyA kit, Ambion AM1350)

1. Denature RNA at 85°C for 2 min or 65°C for 10 min. Cool on ice for 1 min.
2. Set up the reaction by adding :

a. 5X PolyA buffer	10 $\mu$ L
b. 25 mM MnCl <sub>2</sub>	5 $\mu$ L
c. BSA 1mg/mL	2.5 $\mu$ L
d. 100mM ATP (Roche)	1 $\mu$ L
e. Anti- RNase	1 $\mu$ L
f. E_PAP Poly A polymerase	2.5 $\mu$ L
g. RNA	28 $\mu$ L
3. Incubate at 37°C for 20 min.
4. Proceed to small RNA column cleanup, then ethanol precipitation.
5. After drying the pellet, resuspend in 11  $\mu$ L H<sub>2</sub>O. Keep 1  $\mu$ L for running a small RNA Bioanalyzer chip.

#### Ligation

We recently reduced the amount 5'SBS3 adapter by 100 fold. We found that this reduced the amount of linker-linker formed and facilitated its subsequent removal.

1. Denature RNA at 85°C for 2 min or 65°C for 10 min. Cool on ice for 1 min.
2. Set up the reaction by adding :

a. Anti-RNase	0.5 $\mu$ L
b. 10X T4 ligase buffer	1.5 $\mu$ L
c. BSA	0.25 $\mu$ L
d. 5'SBS 3 adapter (1 $\mu$ M)	2 $\mu$ L
e. T4 RNA ligase (Ambion 5U/ $\mu$ L)	0.75 $\mu$ L
f. RNA	10 $\mu$ L
3. Incubate at 4 °C overnight.
4. Proceed to small RNA column cleanup, then ethanol precipitation.
5. After drying the pellet, resuspend in 16.5  $\mu$ L H<sub>2</sub>O.

#### First Strand cDNA Synthesis

1. To 16.5  $\mu$ L RNA, add 2  $\mu$ L A-Tail RT primer.
2. Incubate at 65 °C for 5 min. Leave on ice for 5 min.
3. Then add the following :

a. 10mM dNTPs	1.5 $\mu$ L
b. 5X first strand cDNA buffer	6 $\mu$ L
c. 0.1M DTT	1.5 $\mu$ L
d. Anti-RNase	1 $\mu$ L
e. Superscript RT III	1.5 $\mu$ L

#### PCR

1. Set up the reaction by adding :

a. First strand cDNA	5 $\mu$ L
b. PE 5' PCR primer 100 $\mu$ M	0.5 $\mu$ L
c. PE 3' PCR primer 100 $\mu$ M	0.5 $\mu$ L
d. 2X Phusion mix (NEB F-531L)	50 $\mu$ L
e. H <sub>2</sub> O	44 $\mu$ L
2. Program the thermal cyclers as follows:

1. 94°C      2 min.
  2. 94°C      15 s.
  3. 54 °C     30 s.
  4. 72 °C     20 s.
  5. Go back to step 2 and repeat 4 more times.
  6. 94°C      15 s.
  7. 60 °C     30 s.
  8. 72 °C     20 s.
  9. Go back to step 6 and repeat 12 more times.
  10. 4 °C forever
3. After PCR, clean up the reaction by putting through a Minelute column as follows:
- a. Add 5 volumes of PB buffer to the reaction.
  - b. To bind DNA, apply the sample to a Minelute column. Spin at 13000rpm for 1 min. Discard the flow-through.
  - c. Wash with 750 µL PE buffer. Spin at 13000rpm for 1 min. Discard the flow-through.
  - d. Spin at 13000 rpm for 1 min to dry the column.
  - e. Add 10 µL EB buffer. Spin at 13000rpm for 1 min to elute DNA.
  - f. Repeat the elution one more time.
  - g. Use 1 µL for running on a High sensitivity DNA Bioanalyzer chip.

### Gel extraction

1. Run the rest of the sample in a 2% agarose gel.
2. Excise the DNA from >134bp to 350bp.
3. Weigh the gel slice and add 3 volumes of QG buffer.
4. Incubate at 50°C for 10 min or until the gel slice has completely dissolved.
5. After the gel slice has dissolved completely, check that the color of the mixture is yellow. **Note:** If the color of the mixture is orange or violet, add 10 µL of 3M NaOAc pH 5.0 and mix. The color of the mixture will turn to yellow.
6. Add 1 gel volume of isopropanol to the sample and mix by inverting the tube several times.
7. Place a Minelute column in a provided 2 mL collection tube.
8. To bind DNA, apply the sample to the Minelute column and spin for 1 min. The maximum volume of the column reservoir is 800 µL. For sample volumes of more than 800 µL, simply load and spin again.
9. Discard the flow-through and place the Minelute column back in the same collection tube.
10. Add 500 µL QG buffer to the spin column and spin for 1 min.
11. Discard the flow-through and place the Minelute column back in the same collection tube.
12. To wash, add 750 µL PE buffer to the Minelute column and spin for 1 min.
13. Discard the flow-through and spin the column for an additional 1 min at >10,000g.
14. Place the Minelute column into a clean 1.5 mL tube.
15. To elute DNA, add 10 µL EB buffer to the center of the membrane, let the column stand for 1 min. and spin for 1 min.
16. Repeat the elution to get a higher yield of DNA.
17. Proceed to ethanol precipitation.

### Quantification

1. After drying the pellet, resuspend in 20 µL H<sub>2</sub>O.
2. Measure the concentration on the Nanodrop.
3. Run 1 uL on the High sensitivity DNA Bioanalyzer chip.





